

Induction of Labor EHR Templates

SmartPhrase and order set examples to support standardized induction of labor (IOL) documentation in the electronic health record (EHR).

SmartPhrases

Some clinicians use SmartPhrases as a guide to remember what to discuss and therefore require a descriptive body. Other clinicians prefer 1- 2 sentences of concise language. These example SmartPhrases are meant to be modified to fit the needs of your clinicians and site-specific workflow.

Dual Ripening Agent Recommended

.IOLDUAL

Date/Time

Dual agent ripening {specify option}

- Yes - misoprostol and balloon recommended and accepted
- No - misoprostol recommended, balloon not feasible
 - If this option, current ripening: free text
- Yes - oxytocin and balloon recommended and accepted
- No - oxytocin recommended, balloon not feasible
 - If this option, current ripening: free text
- No - recommended, patient declined dual agent
 - If this option, current ripening: free text

Amniotomy Recommended

.AMNIOTOMY_RECOMMENDED

Amniotomy Recommended Documentation

- Date/Time Recommended: {DATE/TIME}
- Amniotomy Recommended: Yes
- Outcome:
- Patient declined amniotomy after discussion
- Amniotomy not performed due to medical reason:
- Fetal station at -3 or higher
- Fetal head ballotable
- Fetal head floating
- Other indication documented: {specify}
- Plan: Will readdress and consider amniotomy at next cervical exam

Order Sets

The below order sets are meant to be a starting place for creating a tailored order set for your own hospital.

Sample Misoprostol Order Set

misoPROStol (CYTOTEC) tablet 25 mcg:

25 mcg, Buccal, EVERY 3 HOURS PRN, cervical ripening, 6 doses

Starting on MM/DD/YY at TIME, Until Discontinued

misoPROStol (CYTOTEC) tablet 25 mcg:

25 mcg, Vaginal, EVERY 3 HOURS PRN, cervical ripening, 6 doses

Starting on MM/DD/YY at TIME, Until Discontinued

Foley Balloon Order Set

Frequency: Starting on MM/DD/YY at TIME, Until Discontinued

Notify Clinician: If patient experiences a drop in blood pressure, dizziness, nausea, sweating, vaginal bleeding other than bloody show or uterine tachysystole

Notify Clinician: If the device is spontaneously expelled or remains in place 12 hours after placement